

WPAS - Factor List
WPAS Handicap System
Season 2019-2020



Versions: SL: 105 GS: 111 SG: 76 DH: 40

August 2019

Class		SL	GS	SG	DH
B 1	Blind Athlete	0.5647	0.6043	0.6059	0.6132
B 2	Visually Impaired Athlete with a visual acuity of 2/60	0.8084	0.8645	0.8656	0.8760
B 3	Visually Impaired Athlete with a visual acuity above 2/60 to 6/60	0.8453	0.8912	0.8941	0.9182
LW 1	Double above knee amputation	0.8207	0.8380	0.8384	0.8588
LW 2	Single above knee amputation	0.9704	0.8782	0.9077	0.9222
LW 3	Double below knee amputation, minimal through the ankle joint. CP 5 - moderate to slight diplegic involvement CP 6 - moderate athetoid or ataxic impairment	0.8741	0.9080	0.9249	0.9402
LW 4	Single below knee amputation	0.9628	0.9713	0.9795	0.9868
LW 5/7-1	Double amputation of upper limbs above elbow. Motor paresis or paralysis or dysmelia of both upper limbs comparable to 5/7 1 a.	0.9440	0.9545	0.9720	0.9749
LW 5/7-2	a. One arm above elbow and one arm below elbow amputation. b. Motor paresis or paralysis or dysmelia of both upper limbs comparable to 5/7 2 a	0.9478	0.9583	0.9739	0.9758
LW 5/7-3	a. Double amputation of upper limbs below elbow. b. Motor paresis or paralysis or dysmelia of both upper limbs comparable to 5/7 3 a	0.9526	0.9631	0.9788	0.9807
LW 6/8-1	a. Amputation of one upper limb above elbow. b. Dysmelia of one upper limb, comparable to 6/8 1 a. c. Complete paralysis of one upper limb, with fixation to trunk	0.9660	0.9651	0.9820	0.9860
LW 6/8-2	a. Amputation of one upper limb below elbow. b. Dysmelia of one upper limb comparable to 6/8 2 a. c. Paralysis of one upper limb without fixation to trunk.	1.0000	1.0000	1.0000	1.0000
LW 9-1	Those disabled in one upper limb and one lower limb with above knee amputation or comparable disability (CP 7)	0.8224	0.8639	0.8817	0.8913
LW 9-2	Those disabled in one upper limb and below knee amputation or comparable disability (CP 7)	0.8838	0.9235	0.9363	0.9497
LW 10-1	Athletes with disabilities in the lower limbs, no upper abdominal function and no functional sitting balance. This is normally due to a spinal cord lesion around the level T 5/6.	0.6774	0.7719	0.8243	0.8543
LW 10-2	Athletes with disabilities in the lower limbs, some upper abdominal function and no functional sitting balance. This is normally due to a spinal cord lesion at the levels of T 7-T10	0.7299	0.7977	0.8350	0.8593
LW 11	Sit-ski, disabilities in lower limbs, fair sitting	0.7420	0.8066	0.8460	0.8670
LW 12-1	Athletes with a spinal cord lesion or other disabilities with some function in the lower limbs and a good sitting balance	0.7630	0.8319	0.8654	0.8794
LW 12-2	Athletes with a amputation in the lower limbs	0.7909	0.8370	0.8667	0.8802