

WPAS - Factor List
WPAS Handicap System
Season 2021-2022



Version: HM-04

April 2020

Class	SL	GS	SG	DH
B 1 Blind Athlete	0.5651	0.6053	0.6085	0.6053
B 2 Visually Impaired Athlete with a visual acuity of 2/60	0.8090	0.8671	0.8724	0.8709
B 3 Visually Impaired Athlete with a visual acuity above 2/60 to 6/60	0.8466	0.8947	0.8985	0.9142
LW 1 Double above knee amputation	0.8210	0.8387	0.8364	0.8561
LW 2 Single above knee amputation	0.9706	0.8788	0.9069	0.9218
LW 3 Double below knee amputation, minimal through the ankle joint. CP 5 - moderate to slight diplegic involvement CP 6 - moderate athetoid or ataxic impairment	0.8762	0.9111	0.9232	0.9421
LW 4 Single below knee amputation	0.9622	0.9718	0.9801	0.9852
LW 5/7-1 Double amputation of upper limbs above elbow. Motor paresis or paralysis or dysmelia of both upper limbs comparable to 5/7 1 a.	0.9453	0.9515	0.9695	0.9681
LW 5/7-2 a. One arm above elbow and one arm below elbow amputation. b. Motor paresis or paralysis or dysmelia of both upper limbs comparable to 5/7 2 a	0.9491	0.9553	0.9714	0.9691
LW 5/7-3 a. Double amputation of upper limbs below elbow. b. Motor paresis or paralysis or dysmelia of both upper limbs comparable to 5/7 3 a	0.9539	0.9601	0.9763	0.9740
LW 6/8-1 a. Amputation of one upper limb above elbow. b. Dysmelia of one upper limb, comparable to 6/8 1 a. c. Complete paralysis of one upper limb, with fixation to trunk	0.9643	0.9653	0.9805	0.9834
LW 6/8-2 a. Amputation of one upper limb below elbow. b. Dysmelia of one upper limb comparable to 6/8 2 a. c. Paralysis of one upper limb without fixation to trunk.	1.0000	1.0000	1.0000	1.0000
LW 9-1 Those disabled in one upper limb and one lower limb with above knee amputation or comparable disability (CP 7)	0.8226	0.8646	0.8821	0.8892
LW 9-2 Those disabled in one upper limb and below knee amputation or comparable disability (CP 7)	0.8832	0.9239	0.9348	0.9483
LW 10-1 Athletes with disabilities in the lower limbs, no upper abdominal function and no functional sitting balance. This is normally due to a spinal cord lesion around the level T 5/6.	0.6782	0.7756	0.8250	0.8509
LW 10-2 Athletes with disabilities in the lower limbs, some upper abdominal function and no functional sitting balance. This is normally due to a spinal cord lesion at the levels of T 7-T10	0.7306	0.8025	0.8389	0.8586
LW 11 Sit-ski, disabilities in lower limbs, fair sitting	0.7405	0.8110	0.8488	0.8658
LW 12-1 Athletes with a spinal cord lesion or other disabilities with some function in the lower limbs and a good sitting balance	0.7613	0.8356	0.8679	0.8766
LW 12-2 Athletes with a amputation in the lower limbs	0.7913	0.8413	0.8687	0.8774